

# Persistent homology of groups

Graham Ellis & Simon King\*

School of Mathematics, National University of Ireland, Galway

September 1, 2010

## Abstract

We introduce and investigate notions of persistent homology for  $p$ -groups and for coclass trees of  $p$ -groups. Using computer techniques we show that persistent homology provides fairly strong homological invariants for  $p$ -groups of order  $\leq 81$ . The strength of these invariants, together with some of their elementary theoretical properties, suggest that persistent homology may be a useful tool in the study of prime-power groups. In particular, we ask whether the known periodic structure on coclass trees is reflected in a periodic structure on the persistent homology of  $p$ -groups in the trees.

## 1 Introduction

Persistent homology is a tool from applied topology that was introduced for studying qualitative properties of large empirical data sets [4]. At its simplest, the idea is to impose some metric on a data set  $X$ , choose an appropriate sequence of metric space inclusions  $X = X_0 \subset X_1 \subset X_2 \subset \cdots \subset X_N$ , and then study the behaviour of the induced homology maps  $H_n(X_0, \mathbb{F}) \rightarrow H_n(X_1, \mathbb{F}) \rightarrow H_n(X_2, \mathbb{F}) \rightarrow \cdots \rightarrow H_n(X_N, \mathbb{F})$  in a given degree  $n$ . The coefficients  $\mathbb{F}$  are typically chosen to be a field. Such a sequence of linear maps is then determined, up to isomorphism, by an upper triangular matrix  $P_n = (p_{i,j})$  with  $p_{i,j}$  the dimension of the image of the map  $H_n(X_i, \mathbb{F}) \rightarrow H_n(X_j, \mathbb{F})$ . In particular  $p_{i,i}$  is the dimension of  $H_n(X_i, \mathbb{F})$ . The matrix  $P_n$  contains information on the extent to which homology  $n$ -cycles persist through lengths of the induced sequence. Cycles that persist for a significant length are deemed to be significant and, for appropriately chosen  $X_i$ , are likely to represent some qualitative feature of the initial data set  $X$ . Cycles that persist for only a short length are deemed to be less significant.

Persistent homology analysis can be applied to any data set  $X$  for which we have a suitable topology, and for which we have a meaningful sequence of topological inclusions. It provides a concise set of numerical descriptors for homological features

---

\*Supported by Marie Curie fellowship MTKD-CT-2006-042685

of (the sequence of inclusions associated to)  $X$ . In this paper we investigate the potential of applying the idea to groups and particularly to finite  $p$ -groups.

We can view the elements of a group  $G$  as being the vertices of a Cayley graph. Furthermore, we can view the Cayley graph as the 1-skeleton of the universal cover  $EG$  of a classifying CW-space  $BG$ . We set  $X = BG$  and construct an inclusion  $X \hookrightarrow X_1$  from any surjective group homomorphism  $\phi: G \twoheadrightarrow Q$  with  $X_1 = BQ$  a classifying space obtained by attaching cells to  $BG$ . A sequence of inclusions  $X \hookrightarrow X_1 \hookrightarrow \dots \hookrightarrow X_N$  corresponds to a sequence  $G \twoheadrightarrow Q_1 \twoheadrightarrow Q_2 \twoheadrightarrow \dots \twoheadrightarrow Q_N$  of surjective group homomorphisms, or equivalently, to an increasing sequence  $N_1 \leq N_2 \leq \dots \leq N_N$  of normal subgroups of  $G$  (where  $N_i$  is the kernel of the composite surjection  $G \twoheadrightarrow Q_i$ ).

We focus on finite prime-power groups  $G$ , and on the following five normal series in  $G$ .

$$\begin{aligned}
L_1(G) &= G, & L_{i+1}(G) &= [L_i(G), G] && \text{(lower central)} \\
L_1^p(G) &= G, & L_{i+1}^p(G) &= [L_i^p(G), G](L_i^p(G))^p && \text{(lower } p\text{-central)} \\
D_1(G) &= G, & D_{i+1}(G) &= [D_i(G), D_i(G)] && \text{(derived)} \\
Z_0(G) &= 1, & Z_{i+1}(G) &= \text{preimage of } Z(G/Z_i(G)) \text{ in } G && \text{(upper central)} \\
Z_0^p(G) &= 1, & Z_{i+1}^p(G) &= \text{elements of order } \leq p \text{ in the} \\
&&& \text{preimage of } Z(G/Z_i^p(G)) \text{ in } G && \text{(upper } p\text{-central)}
\end{aligned}$$

These five series can be regarded as functors from the category whose objects are groups and whose arrows are surjections of groups. They can be regarded as functors to the category whose objects are sequences of group homomorphisms, and whose morphisms are commutative diagrams of groups. So, for instance, we view  $Z$  as a functor which sends a surjection  $G \rightarrow Q$  to the following commutative diagram.

$$\begin{array}{ccccccc}
G & \rightarrow & G/Z_1(G) & \rightarrow & G/Z_2(G) & \rightarrow & \dots \\
\downarrow & & \downarrow & & \downarrow & & \\
Q & \rightarrow & Q/Z_1(Q) & \rightarrow & Q/Z_2(Q) & \rightarrow & \dots
\end{array}$$

For  $F$  equal to any of  $L, L^p, D, Z, Z^p$  we define the *persistence matrix*  $P_n^F(G) = (p_{i,j})$  to be an upper triangular matrix. For  $F$  equal to  $Z$  or  $Z^p$  and  $i \geq j$  the entry  $p_{i,j}$  is the dimension of the image of the map

$$h_{i,j}: H_n(B(G/F_i(G)), \mathbb{F}_p) \rightarrow H_n(B(G/F_j(G)), \mathbb{F}_p).$$

For  $F$  equal to  $L, L^p$  or  $D$  and  $i \geq j$  the entry  $p_{i,j}$  is the dimension of the image of the map

$$h_{i,j}: H_n(B(G/F_j(G)), \mathbb{F}_p) \rightarrow H_n(B(G/F_i(G)), \mathbb{F}_p).$$

The family  $H_n^F(G) = \{h_{i,j}\}_{i \geq j}$  is called a *persistence module* and is a functorial invariant of the group  $G$ . Two persistence modules are isomorphic if and only if the corresponding persistence matrices are identical.

Our aim is to investigate the extent to which persistence matrices can be used to capture the structure of finite  $p$ -groups. For instance, we derive some elementary theoretical results aimed at understanding the nature of the group-theoretic information contained in persistence matrices: Even the simplest terms of the persistence

data yield useful information. In addition we study, by means of computer techniques [9, 8, 6, 7], the strength of persistent homology as an invariant of groups. We establish that the degree seven upper central series persistence matrix  $P_7^Z(G)$  yields 181 distinct matrices when  $G$  ranges over the 267 groups of order 64. Furthermore, the groups of order 64 give rise to 187 distinct infinite sequences  $P_*^Z(G) = (P_n^Z(G))_{n \geq 1}$ . We give analogous statistics for each of the five series  $F$  and all prime-power groups of order at most 81. We believe that the theoretical properties of persistence matrices, in conjunction with their apparent strength as group invariants, suggest that persistent homology may be a convenient tool in the study of prime-power groups. In particular, we end the paper with the question of whether the known periodic structure on coclass trees is somehow reflected in a periodic structure on the persistent homology of  $p$ -groups in the trees.

## 2 Examples and properties of persistence

Consider the dihedral group  $D_{32}$  of order 64. Using algorithms recently implemented in the group cohomology package [9] for SAGE (see [8] for an overview of its algorithms) or the GAP homological algebra package HAP [6] (see [7] for an overview of its algorithms) one can compute the lower central series persistence matrix of  $D_{32}$  in degree 2 to be

$$P_2^L(D_{32}) = \begin{pmatrix} 3 & 2 & 2 & 2 & 2 \\ 0 & 3 & 2 & 2 & 2 \\ 0 & 0 & 3 & 2 & 2 \\ 0 & 0 & 0 & 3 & 2 \\ 0 & 0 & 0 & 0 & 3 \end{pmatrix}.$$

The first row of this matrix implies that  $H_2(D_{32}, \mathbb{F}_2)$  has dimension 3, and that precisely two basis elements persist (*i.e.* remain non-zero) under the induced maps  $H_2(D_{32}, \mathbb{F}_2) \rightarrow H_2(D_{32}/L_i(D_{32}), \mathbb{F}_2)$  ( $2 \leq i \leq 5$ ). The second row implies that  $H_2(D_{32}/L_5(D_{32}), \mathbb{F}_2)$  has a basis of three elements, precisely two of which persist under the induced maps  $H_2(D_{32}/L_5(D_{32}), \mathbb{F}_2) \rightarrow H_2(D_{32}/L_i(D_{32}), \mathbb{F}_2)$  ( $2 \leq i \leq 4$ ). The matrix is represented by the *persistence bar code* shown in Figure 1(b) and, in fact, can be reconstructed from the information in this bar code.

We regard a persistence bar code as a directed graph whose vertices are arranged in columns, whose edges connect certain pairs of vertices in neighbouring columns, and where all edges are horizontal and directed away from the first column. The use of bar codes for describing persistence matrices was introduced by G. Carlsson *et al.* in [2].

Persistence bar codes for the matrices  $P_n^L(D_{32})$ ,  $n = 1, 3$ , are given in Figures 1(a) and (c). Persistence bar codes for the matrices  $P_n^L(Q_{32})$ ,  $n = 1, 2, 3$ , associated to the quaternion group of order 64 are given in Figure 2. Persistence bar codes for the matrices  $P_n^L(QD_{32})$ ,  $n = 1, 2, 3$ , associated to the quasi-dihedral group of order 64 are given in Figure 3. (Recall that for every  $n \geq 4$  there are exactly four isomorphism types of nonabelian groups of order  $2^n$  with a cyclic subgroup of index 2. Precisely three of these have nilpotency class  $n - 1$ , namely the dihedral group, generalized quaternion group and the *quasi-dihedral* group.)




Figure 1: Degree 1, 2 and 3 lower central bar codes for  $D_{32}$




Figure 2: Degree 1, 2 and 3 lower central bar codes for  $Q_{32}$




Figure 3: Degree 1, 2 and 3 lower central bar codes for  $QD_{32}$

We are interested in the extent to which group-theoretic information is retained by persistence matrices for the functors  $L$ ,  $L^p$ ,  $D$ ,  $Z$ ,  $Z^p$ . We shall let  $F$  denote any one of the above five functors and let  $F_i(G)$  denote the  $i$ th term in the corresponding normal subseries of  $G$ .

**Proposition 1** *Let  $G$  be a  $p$ -group.*

(i) *For  $F$  any of the above five functors the first persistence matrix  $P_1^F(G)$  determines the minimal number of generators of  $G/F_i(G)$  for all  $i \geq 1$ .*

(ii) *For  $F = L$  or  $Z$  the first persistence matrix  $P_1^F(G)$  determines the nilpotency class of  $G$ .*

(iii) *For  $F = L^p$  or  $Z^p$  the first two persistence matrices  $P_1^F(G)$  and  $P_2^F(G)$  determine the order of  $G$ .*

**Proof.** Assertion (i) follows from the fact that the dimension of the vector space  $H_1(G/F_i(G), \mathbb{F}_p) \cong G/F_i(G)[G, G]G^p$  is equal to the minimal number of generators of  $G/F_i(G)$ . This dimension is the entry  $p_{i,i}$  in the first persistence matrix. Assertion (ii) is just the observation that the number of columns in the persistence matrix is by definition equal to the length of the upper or lower central series of  $G$ . We prove assertion (iii) just for the functor  $F = Z^p$ . We use induction on the length  $k$  of the upper  $p$ -central series. If  $k=1$  then  $G$  is the trivial group. If  $k = 2$  then  $G$  is an elementary abelian  $p$ -group of order  $p^d$  where  $d$  can be determined by (i). As an inductive hypothesis suppose that the assertion is true when the upper  $p$ -central series has length  $k$ . For  $G$  a group with upper  $p$ -central series of length  $k + 1$  we set  $Q = G/Z_1^p(G)$ . The five term natural exact homology sequence

$$H_2(G, \mathbb{F}_p) \rightarrow H_2(Q, \mathbb{F}_p) \rightarrow Z_1^p(G) \rightarrow H_1(G, \mathbb{F}_p) \rightarrow H_1(Q, \mathbb{F}_p) \rightarrow 0 \quad (1)$$

allows us to determine the dimension of the vector space  $Z_1^p(G)$  from the first two upper  $p$ -central persistence matrices. By the inductive hypothesis we can determine the order of  $Q$  from these two matrices. Then we have the order  $|G| = |Q||Z_1^p(G)|$  as required.  $\square$

**Proposition 2** *The abelian invariants of an abelian  $p$ -group  $G$  are determined by the first upper  $p$ -central persistence matrix  $P_1^{Z^p}(G)$ .*

**Proof.** Suppose that

$$G = C_{p^{n_1}} \times C_{p^{n_2}} \times \cdots \times C_{p^{n_d}}$$

with each  $n_i \geq 1$ . It suffices to observe that the bar code for  $P_1^{Z^p}(G)$  consists of  $d$  horizontal bars, each starting in the first column, the  $i$ th bar reaching as far as the  $n_i$ th column. (As an example, this bar code is given in Figure 4 for the group  $G = C_2 \times C_4 \times C_4 \times C_{16}$ .) The observation can be proved by induction on the length  $k$  of the upper  $p$ -central series. If  $k = 1$  then the persistence matrix has just one entry, namely the dimension of the elementary abelian group  $G$ . In general we set  $Q = G/Z_1^p(G)$  and assume our description of the bar code for  $P_1^{Z^p}(Q)$ . The surjection  $G \rightarrow Q$  of abelian groups induces a surjection in first homology  $H_1(G, \mathbb{F}_p) \rightarrow H_1(Q, \mathbb{F}_p)$ . The bar code for  $P_1^{Z^p}(G)$  is thus obtained from that for




Figure 4: First upper 2-central bar code for  $G = C_2 \times C_4 \times C_4 \times C_{16}$

$P_1^{Zp}(Q)$  by adding a new first column of  $d$  vertices together with horizontal edges connecting each vertex in the new second column to some vertex in the new first column.  $\square$

**Proposition 3** *Let  $G$  be a finite  $p$ -group.*

(i) *The bar code for the first lower central persistence matrix  $P_1^L(G)$  consists of  $d = \text{rank}(G/[G, G]G^p)$  horizontal paths, each starting in the first column and ending in the final column.*

(ii) *In the bar code for the second lower central persistence matrix  $P_2^L(G)$  every horizontal path starts in the first column.*

(iii) *In the bar code for the second lower central persistence matrix  $P_2^L(G)$  let  $v$  be the number of vertices in the  $j$ th column ( $j \geq 2$ ) that are not incident with an edge. Set  $j' = c+2-j$  where  $c$  is the nilpotency class of  $G$ . Then  $v$  is the dimension of the vector space  $L_{j'}(G)/L_{j'+1}(G) \otimes \mathbb{F}_p$ . (In particular, the number of vertices in the right-most column not incident with an edge is equal to the rank of  $L_2(G)/L_3(G) \otimes \mathbb{F}_p$ .)*

**Proof.** Assertion (i) follows from the canonical isomorphisms  $H_1(G/L_i(G), \mathbb{F}_p) \cong G/[G, G]G^p$  for  $i \geq 0$ . Assertion (ii) follows from the natural exact sequences

$$H_2(G/L_{i+1}(G), \mathbb{F}_p) \rightarrow H_2(G/L_i(G), \mathbb{F}_p) \rightarrow L_i(G)/L_{i+1}(G) \otimes \mathbb{F}_p \rightarrow 0$$

$$H_2(G, \mathbb{F}_p) \rightarrow H_2(G/L_i(G), \mathbb{F}_p) \rightarrow L_i(G)/L_{i+1}(G) \otimes \mathbb{F}_p \rightarrow 0$$

that arise as part of the five term exact sequence in mod- $p$  homology. These two sequences imply that the two homomorphisms  $H_2(G/L_{i+1}(G), \mathbb{F}_p) \rightarrow H_2(G/L_i(G), \mathbb{F}_p)$  and  $H_2(G, \mathbb{F}_p) \rightarrow H_2(G/L_i(G), \mathbb{F}_p)$  have identical images. This identity implies (ii). Assertion (iii) follows from the second of the exact sequences.  $\square$

### 3 Persistence matrices as group invariants

Proposition 2 states that  $P_1^{Zp}$  is a complete invariant for abelian  $p$ -groups. We now investigate the strength of persistent homology as a group invariant for finite prime-power groups of low order. The computations were made using the second author's group cohomology package [9] for the computer algebra system Sage [11]. Where possible, the computations were corroborated using the first author's GAP package [6]). We begin with the following summary of the computations for  $P_*^F(G) = (P_n^F(G))_{n \geq 1}$ .

$F$		$k = 8$	$k = 16$	$k = 27$	$k = 32$	$k = 64$	$k = 81$
	$\text{Nr}(k)$	5	14	5	51	267	15
$Z$	$( C , \max)$	(5, 1)	(13, 2)	(5, 1)	(44, 2)	(187, 7)	(14, 2)
	$t$	3	4	3	5	6	5
	$( C , \max, d)$	(5, 1, 3)	(13, 2, 4)	(5, 1, 3)	(44, 2, 7)	(181, 7, 7)	(14, 2, 5)
$Z^p$	$( C , \max)$	(5, 1)	(13, 2)	(5, 1)	(42, 3)	(174, 8)	(14, 2)
	$t$	3	4	3	5	6	5
	$( C , \max, d)$	(5, 1, 3)	(13, 2, 4)	(5, 1, 3)	(42, 3, 5)	(166, 8, 7)	(14, 2, 5)

Table 1:

**Theorem 4** For the 366 groups of order at most 81:

- (i)  $P_*^Z$  partitions the groups into 277 classes with maximum class size equal to 7.
- (ii)  $P_*^{Z^p}$  partitions the groups into 262 classes with maximum class size equal to 8.
- (iii)  $P_*^L$  partitions the groups into 227 classes with maximum class size equal to 7.
- (iv)  $P_*^{L^p}$  partitions the groups into 179 classes with maximum class size equal to 15.
- (v)  $P_*^D$  partitions the groups into 180 classes with maximum class size equal to 13.

A more detailed description of the computations is given in Tables 1 and 2 which contain, for prime-powers  $k = 8, 16, 27, 32, 64, 81$ , and for each of the five functors  $F$ :

1. the number  $\text{Nr}(k)$  of isomorphism classes of groups of order  $k$ .
2. the integer pair  $(|C|, \max)$  where  $|C|$  is the number of classes of groups of order  $k$  distinguished by the invariant  $P_*^F$ , and  $\max$  is the maximum size of a class.
3. the smallest integer  $t$  for which the invariant  $P_{*\leq t}^F = (P_n^F)_{n\leq t}$  is as strong as  $P_*^F$  on the groups of order  $k$ .
4. an integer triple  $(|C|, \max, d)$  where  $|C|$  is the number of classes of groups of order  $k$  distinguished by the matrix  $P_d^F$ , and  $\max$  is the maximum size of a class (for some choice of  $d$ ).

Persistence bar codes can sometimes distinguish between very similar groups. Consider for example the two groups  $G, G'$  of order 64 which are given the identification numbers 158 and 160 in the Small Groups Library [1] that is available in GAP. Their mod-2 cohomology rings  $H^*(G, \mathbb{F}_2)$  and  $H^*(G', \mathbb{F}_2)$  have: the same Poincaré

$F$		$k = 8$	$k = 16$	$k = 27$	$k = 32$	$k = 64$	$k = 81$
	$\text{Nr}(k)$	5	14	5	51	267	15
$L$	$( C , \max)$	(5, 1)	(12, 2)	(5, 1)	(37, 3)	(145, 7)	(14, 2)
	$t$	3	5	3	5	6	5
	$( C , \max, d)$	(5, 1, 3)	(12, 2, 4)	(5, 1, 3)	(37, 3, 5)	(144, 7, 9)	(14, 2, 5)
$L^p$	$( C , \max)$	(4, 2)	(9, 2)	(5, 1)	(28, 5)	(110, 15)	(14, 2)
	$t$	3	4	3	5	6	5
	$( C , \max, d)$	(4, 2, 3)	(9, 2, 4)	(5, 1, 3)	(28, 5, 5)	(109, 15, 9)	(14, 2, 5)
$D$	$( C , \max)$	(5, 1)	(10, 2)	(5, 1)	(29, 5)	(108, 13)	(14, 2)
	$t$	3	4	3	5	6	5
	$( C , \max, d)$	(5, 1, 3)	(10, 2, 4)	(5, 1, 3)	(29, 5, 7)	(106, 13, 11)	(14, 2, 5)

Table 2:

		$k = 8$	$k = 16$	$k = 27$	$k = 32$	$k = 64$	$k = 81$
	$\text{Nr}(k)$	5	14	5	51	267	15
$H_{*\leq 3}^{Z^p}$	$( C , \max)$	(5, 1)	(13, 2)	(5, 1)	(46, 3)	(188, 8)	(14, 2)

Table 3:



Figure 5: Degree 3 upper central bar codes for groups 158 and 160 of order 64.

series; the same number of generators and relations sorted by degree; the same depth; the same  $a$ -invariants. Both groups have nilpotency class 3 and their upper central series admit isomorphisms  $Z_n(G) \cong Z_n(G')$ ,  $Z_{n+1}(G)/Z_n(G) \cong Z_{n+1}(G')/Z_n(G')$  for  $1 \leq n \leq 3$ . Their  $p$ -upper central series, lower central series,  $p$ -lower central series and derived series admit analogous isomorphisms. However, their upper central bar codes shown are different in degree 3 (see Figure 5).

## 4 Integral persistence

One can use integral homology groups in place of mod  $p$  homology groups when studying persistence. However, the induced homology homomorphisms  $H_n(f): H_n(G, \mathbb{Z}) \rightarrow H_n(Q, \mathbb{Z})$  can not be fully described using the notion of dimension. A partial description is given by the abelian invariants of the source, the target and the cokernel of  $H_n(f)$ . It is partial due to the extension problem. We denote by  $IP_n^F(G)$  the upper triangular matrix whose entry in the  $i$ -th row and  $j$ -th column ( $j \geq i$ ) is a triple  $(A, B, C)$  containing lists of the abelian invariants of the source, target and cokernel of the map  $H_n(G/F_i(G), \mathbb{Z}) \rightarrow H_n(G/F_j(G), \mathbb{Z})$ . Table 3 indicates the strength of  $IP_{* \leq t}^{Z^p} = (IP_n^{Z^p})_{n \leq t}$  as an invariant of the prime-power groups of order at most 81 for  $t = 3$ .

## 5 Persistence in coclass trees

Recall that a  $p$ -group of order  $p^n$  and nilpotency class  $c$  is said to have *coclass*  $r = n - c$ . For a fixed  $p$  and  $r$  one can consider the graph  $\mathbb{G}(p, r)$  whose vertices are the  $p$ -groups of coclass  $c$ ; two groups  $G$  and  $H$  are connected by an edge in the graph if and only if there exists a normal subgroup  $N \leq H$  of order  $p$  such that  $H/N \cong G$ . It is not difficult to observe that the graph  $\mathbb{G}(p, r)$  is a forest with infinitely many vertices.

The graph  $\mathbb{G}(p, r)$  can be stratified into levels by deeming all groups of order  $p^l$  to be at level  $l$ . The graph  $\mathbb{G}(2, 1)$  is shown in Figure 6. Its three columns contain, respectively, the dihedral groups of order  $2^l$ , the quaternion groups of order  $2^l$  and the quasi-dihedral groups of order  $2^l$ . Lower central bar codes for the three coclass 1 groups of order 128 are shown in Figures 1, 2, 3.



The philosophy is that  $PH_n(\mathbb{T})$  should capture some group-theoretic properties that are common to all groups in the tree. Easy results in this direction are parts (ii) and (iv) of the following proposition. Part (i) of the proposition implies that the surjections in (2) are isomorphisms for all sufficiently large  $l$ . Hence  $PH_n(\mathbb{T}) = \text{Image}(H_n(G_{l+1}, \mathbb{F}_p) \rightarrow H_n(G_l, \mathbb{F}_p))$  for all groups  $G_l$  on the infinite path in the tree above a certain level.

**Proposition 7** (i) *The persistent homology  $PH_n(\mathbb{T})$  is a finite dimensional vector space for all degrees  $n \geq 1$ .*

(ii) *The dimension of  $PH_1(\mathbb{T})$  equals the minimum number of generators for any group in the tree.*

(iii)  *$H_2(G, \mathbb{F}_p) \cong PH_2(\mathbb{T}) \oplus \mathbb{F}_p$  for all groups  $G$  above a certain level in the tree which are not leaves. For leaves, the dimension of  $H_2(G, \mathbb{F}_p)$  is at least that of  $PH_2(\mathbb{T})$ .*

(iv) *Any group  $G$  in the tree needs at least  $\dim(PH_2(\mathbb{T}))$  relators to define it. If the group is not a leaf it needs at least  $\dim(PH_2(\mathbb{T})) + 1$  relators.*

**Proof.** (i) The  $p$ -adic space group associated to  $\mathbb{T}$  decomposes into a short exact sequence  $1 \rightarrow T \rightarrow S_{\mathbb{T}} \rightarrow P \rightarrow 1$  where  $P$  is a finite (point) group. Each group  $G_l$  on the infinite path of  $\mathbb{T}$  thus fits into a short exact sequence  $1 \rightarrow T/T_l \rightarrow G_l \rightarrow P \rightarrow 1$ . Let  $R_*^P \rightarrow \mathbb{Z}$  be any free  $\mathbb{Z}P$ -resolution of the integers. Let  $R_*^{(T/T_l)} \rightarrow \mathbb{Z}$  be the minimal free  $\mathbb{Z}(T/T_l)$ -resolution of the integers constructed as a tensor product of resolutions of the cyclic summands of  $T/T_l$ . Note that the number of free generators of  $R_*^{(T/T_l)}$  in a given degree is independent of  $l$ . By a Lemma of C.T.C. Wall the boundary map in the tensor product  $R_*^{(T/T_l)} \otimes R_*^P$  can be perturbed to produce a free  $\mathbb{Z}G_l$ -resolution  $R_*^{(T/T_l)} \tilde{\otimes} R_*^P$ . By construction, the number of generators of this latter resolution, in a given degree, is independent of  $l$ . This implies that for a given  $n$  the dimension of the homology groups  $H_n(G_l, \mathbb{F}_p)$  is bounded by a number depending only on  $P$  and  $T$ . This means that the sequence of dimensions of the vector spaces  $P_l H_n(\mathbb{T}), P_{l+1} H_n(\mathbb{T}), \dots$  is bounded above. The sequence is also monotonically increasing since the maps in (2) are surjective. Hence the sequence of dimensions converges to the dimension of the inverse limit.

(ii) This follows directly from the definition of  $PH_1(\mathbb{T})$  and the isomorphism  $H_1(G, \mathbb{F}_p) \cong G/[G, G]G^p$ .

(iii) Let  $G_{l+1} \rightarrow G_l$  be a homomorphism in the tree from level  $l+1$  to level  $l$  with kernel  $K$  of order  $p$ . The five term natural exact homology sequence

$$H_2(G_{l+1}, \mathbb{F}_p) \rightarrow H_2(G_l, \mathbb{F}_p) \rightarrow K \rightarrow H_1(G_{l+1}, \mathbb{F}_p) \xrightarrow{\cong} H_1(G_l, \mathbb{F}_p) \rightarrow 0 \quad (3)$$

implies  $\text{Image}(H_n(G_{l+1}, \mathbb{F}_p) \rightarrow H_n(G_l, \mathbb{F}_p)) \oplus K \cong H_2(G_l, \mathbb{F}_p)$ . If  $G_{l+1}$  happens to be on the infinite path in the tree then, by (i), the first term in this sum stabilizes to  $PH_2(\mathbb{T})$  for large  $l$ . If  $G_{l+1}$  is not on the infinite path but is not a leaf, then (i) together with Proposition 3(ii) show that first term in the sum stabilizes to  $PH_2(\mathbb{T})$ . If  $G$  happens to be a leaf then we can at least conclude from Proposition 3 that  $H_2(G_{l+1}, \mathbb{F}_p)$  maps onto  $PH_2(\mathbb{T})$ .

(iv) A presentation for  $G$  yields the low-dimensional terms in a free  $\mathbb{Z}G$ -resolution  $R_*^G \mathbb{Z}$  where the  $\mathbb{Z}G$ -rank of  $R_2^G$  equals the number of relators in the presentation.

Clearly this rank has to be at least the dimension of  $H_2(G, \mathbb{F}_p)$ . So (iii) gives the required result.  $\square$

Persistent homology could easily be computed for some coclass trees. For instance, computer calculations strongly suggest the following result, whose proof should be just a routine homological calculation.

**Conjecture 8** *For  $\mathbb{T}$  the infinite tree in  $\mathbb{G}(2, 1)$  we have*

$$PH_n(\mathbb{T}) = \mathbb{F}_2 \oplus \mathbb{F}_2 \quad (n \geq 1).$$

We note that Proposition 7(iii) is related to Theorem 2 in [5]. The groups in any infinite coclass tree can be canonically partitioned into infinite families. (For instance, the infinite tree in  $\mathbb{G}(2, 1)$  contains three infinite families: the dihedral groups, the quaternion groups and the quasi-dihedral groups.) Theorem 2 in [5] implies that  $H_2(G, \mathbb{F})$  takes on a single value for almost all groups  $G$  in an infinite family. Eick, Feichtenschlager [5] and others have conjectured that the periodic structure of coclass trees is preserved by the functor which maps a group  $G$  to the graded commutative ring  $H^*(G, \mathbb{F})$ . Theorem 2 in [5] confirms the conjecture in degree 2. Additional experimental evidence in support of the conjecture is provided in [3].

We conclude the paper by noting that (limited) experimental evidence suggests that the periodicity in coclass trees might be reflected in a periodic structure on bar codes for the persistence matrices  $P_n^L(G)$  for the groups  $G$  at a sufficiently high level in the tree. Instead of making this statement precise we illustrate it in Figure 7 for  $n = 3$  and  $G$  the mainline group at level 8 in an infinite tree in  $\mathbb{G}(2, 2)$ . The tree and most of the bar code exhibit a periodicity of period 2.

## References

- [1] H.U. Besche, B. Eick & E. O'Brien. A millennium project: constructing small groups. *Internat. J. Algebra Comput.* **12** (2002), 623–644. <http://www-public.tu-bs.de:8080/~hubesche/small.html>
- [2] A. Collins, A. Zomorodian, G. Carlsson & L. Guibas. A barcode shape descriptor for curve point cloud data. *Computers and Graphics* **28**, **6** (2004), 881–894.
- [3] H. Dietrich, B. Eick, and D. Feichtenschlager. Investigating p-groups by coclass with GAP. In *Computational group theory and the theory of groups*, volume 470 of *Contemp. Math.*, pages 4561. Amer. Math. Soc., Providence, RI, 2008.
- [4] H. Edelsbrunner and J. Harer. Persistent homology - a survey. *Contemporary Mathematics* **453** (2008), 257–282.
- [5] B. Eick and D. Feichtenschlager. Computation of low-dimensional (co)homology groups for infinite sequences of p-groups with fixed coclass. Preprint <http://www-public.tu-bs.de:8080/~beick/publ/symbcomp.pdf>.
- [6] G. Ellis. HAP - a homological algebra package for GAP. <http://www.gap-system.org/packages>




Figure 7:  $\mathbb{G}(2, 2)$  tree and third lower central bar code for a mainline group

- [7] G. Ellis. Homological Algebra Programming. *Contemporary Mathematics*, **470** (2008), 63–74.
- [8] D. Green & S. King. The computation of the cohomology rings of all groups of order 128. *J. Algebra*, to appear.
- [9] S. King. Group Cohomology Package. Peer-reviewed optional package for Sage [11] (2009).  
<http://sage.math.washington.edu/home/SimonKing/Cohomology/>
- [10] C. Leedham-Green & S. McKay. *The structure of groups of prime power order*. London Mathematical Society Monographs, New Series 27, Oxford University Press 2002.
- [11] W. A. Stein et al. Sage Mathematics Software, Version 4.2.1 (2009).  
<http://www.sagemath.org>